

Taking Your Hepatitis C Therapy

DACLATASVIR + SOFOSBUVIR with or without Ribavirin

Daclatasvir



- Dose: 1 tablet (usually 60 mg) daily, taken at the same time each day
- You can take with or without food
- Store at room temperature (between 59°F and 86°F)

30 mg ☐
60 mg ☐

Sofosbuvir



- Dose: 1 tablet (400 mg) daily, taken at the same time each day
- You can take with or without food
- Store at room temperature away from direct sunlight

Do You Need to Take Ribavirin? Yes ☐ No ☐



- You will be given 200 mg capsules or tablets; your dose may change during the course of treatment
- Take with food to minimize stomach upset such as nausea and vomiting
- Do not consider pregnancy until at least 6 months after treatment for either partner has ended

My Daily Medication Schedule

Start Date: _____ End Date: _____ Expected Treatment Duration: _____ weeks

Morning	Time: _____ am	Evening	Time: _____ pm	
	Daclatasvir	Take 1 tablet daily		
	Sofosbuvir	Take 1 tablet daily		
Yes <input type="checkbox"/> No <input type="checkbox"/>	Ribavirin	Take _____ ribavirin with food	Ribavirin	Take _____ ribavirin with food

Note: **Daclatasvir** and **sofosbuvir** (and **ribavirin**) can be taken together or at different times, but keep to the same time of day for each drug

If You Miss a Dose:

Daclatasvir

- If you miss a dose, TAKE THE MISSED DOSE THE SAME DAY as soon as you remember; take your next dose of daclatasvir at your regular time the next day
- Do not take more than 1 tablet of daclatasvir in a day
- DO NOT take two doses at the same time to make up for a missed dose

Sofosbuvir

- If you miss a dose, TAKE THE MISSED DOSE THE SAME DAY as soon as you remember; take your next dose of sofosbuvir at your regular time the next day
- Do not take more than 1 tablet of sofosbuvir in a day
- DO NOT take two doses at the same time to make up for a missed dose

Note: If you miss multiple doses of either drug, call your prescriber/clinic listed below

Prescriber/Clinic Contact: _____

For Refills: _____